

## Hepatic Fatty Acids in Wild Rockhopper (*Eudyptes crestatus*) and Magellanic (*Spheniscus magellanicus*) Penguins before and after Moulting

LAURENCE S. HARBIGE<sup>1</sup>, KEB GHEBREMESKEL<sup>2</sup>, GLYNNE WILLIAMS  
and MICHAEL A. CRAWFORD<sup>2</sup>

<sup>1</sup>*Institute of Zoology, Zoological Society of London, Regents Park, London  
NW1 4RY, UK., and* <sup>2</sup>*The Institute of Brain Chemistry and  
Human Nutrition Hackney Hospital, Homerton High  
Street, London E9 6BE, UK.*

**ABSTRACT**—Intra- and inter-species hepatic differences for wild rockhopper (*Eudyptes crestatus*) and magellanic (*Spheniscus magellanicus*) penguin fatty acids were compared both pre-and post-moult. Linoleic (18:2n-6) and arachidonic (20:4n-6) acid composition were significantly higher and palmitic (16:0) acid significantly lower in pre moult rockhopper penguins than in comparable magellanic. Post-moult magellanic had significantly more palmitoleic (16:1), gadoleic (20:1n-9) and erucic, (22:1n-9) and less arachidonic and eicosapentaenoic (20:5n-3) percent fatty acids than post-moult rockhoppers. In both species moulting resulted in a significant reduction in eicosapentaenoic and docosapentaenoic (22:5n-3), and a significant increase in linoleic acid (18:2n-6) percent. In rockhoppers, post moult was associated with an increase in the proportion of palmitic (16:0) and a decrease in palmitoleic (16:1) acid. In the post-moult magellanic, however, there was a decrease in stearic (18:0) and an increase in gadoleic (20:1n-9) and erucic (22:1n-9) fatty acid composition.

### INTRODUCTION

Successful breeding, migration, and moulting in birds are closely linked to nutritional factors [1-4]. Scarcity of food and thus poor nutrient deposition prior to these events can result in failure to breed, poor health and mortality [1-6], especially in species such as penguins which abstain from feeding while breeding and moulting. In penguins moulting lasts between two and five weeks depending on species, during which time body weight losses of 23-60% have been recorded [7-11]. Before breeding and moulting penguins build up body reserves mainly in the form of lipid. These adaptive responses result in body weight increases of 5-33% depending on species and sex [10-12].

Lipids have diverse biological roles; neutral

lipids provide important energy reserves, whilst phospholipids have membrane structural functions [13-15]. The polyunsaturated fatty acid components of phospholipids provide substrates for the cell regulatory molecules the eicosanoids, [14] and are thought to provide structural integrity to cell membranes [16-19].

In view of the importance of lipids, we have investigated the fatty acid composition of hepatic tissue from wild rockhopper (*Eudyptes crestatus*) and magellanic (*Spheniscus magellanicus*) penguins both pre-and post-moult.

### MATERIALS AND METHODS

#### *Penguins*

Liver tissue samples were obtained at necropsy from adult healthy free-living rockhopper and magellanic penguins inhabiting the Falkland Islands during February 1987 after a post-breeding feeding period. This was undertaken as a result of

Accepted October 29, 1991

Received July 17, 1991

<sup>1</sup> Present Address: The Institute of Brain Chemistry and Human Nutrition Hackney Hospital, Homerton High street, London E9 6BE, UK.

a penguin mortality investigation in the Falkland Islands in 1986 [20]. It was not possible to differentiate between adults and subadults even when taking into consideration the appearance of the gonads [11]. The period between the arrival on the moulting area and the beginning of the old feather loss was classified as pre-moult. Whereas, the penguins that had replaced their old feathers with new plumage were considered to be post-moult.

#### Lipid extraction

Total lipids were extracted from liver samples by the method of Folch *et al.*, [21]. Tissues were homogenised in chloroform: methanol (2:1 v/v) containing 0.01% 2,6-di-*t*-butyl-4-methylphenol (BHT) as an antioxidant and left for 24 hr at 4°C. The homogenate-solvent mixture was filtered and transferred to a separatory funnel and left overnight at 4°C following the addition of 25% saline (0.85% NaCl) by volume. The lower organic phase was evaporated in a Rotavap-R (Buchi) under reduced pressure at 37°C. Samples were kept under nitrogen during and after the extraction

procedures and extracts stored at -20°C until required.

#### Fatty acid separation and identification

Total lipids were transmethylated under nitrogen at 70°C for 3 hr with 5 ml of 5% sulphuric acid in methanol as an esterifying reagent. The fatty acid methyl ester derivatives were separated and identified as previously described [22] except that the chromatograph was a Varian model 3700, and the column a CP Sil88 (SP 2340).

#### Statistical analysis

Data are expressed as means and standard deviations with their maximum and minimum ranges. Interspecies mean differences and pre-and post-moult means were compared by Student's unpaired t-test.

## RESULTS

The hepatic fatty acid composition of rockhopper and magellanic penguins pre-and post-moult

TABLE 1. Range and mean  $\pm$  SD percent liver fatty acids (16:0-22:6n-3) in wild pre-and post-moult rockhopper (*Eudyptes crestatus*) and magellanic (*Spheniscus magellanicus*) penguins.

Fatty acid	Rockhopper				Magellanic			
	Pre-moult (n=4)		Post-moult (n=4)		Pre-moult (n=3)		Post-moult (n=4)	
16:0	15.2-19.9	17.9 $\pm 2.1$	21.8-24.1	22.8 $\pm 1.1$	21.9-26.1	24.8 $\pm 2.5$	20.2-24.1	22.9 $\pm 1.8$
16:1	1.3- 3.1	2.3 $\pm 1.0$	0.7- 1.2	0.9 $\pm 0.2$	1.7- 2.1	1.9 $\pm 0.2$	1.7- 3.4	2.6 $\pm 1.1$
18:0	19.2-21.6	20.4 $\pm 1.0$	18.0-23.6	20.8 $\pm 2.9$	18.6-22.8	20.6 $\pm 2.1$	12.4-17.8	15.8 $\pm 2.4$
18:1n-9	18.6-25.3	21.4 $\pm 2.8$	16.5-19.1	18.1 $\pm 1.1$	18.0-22.4	20.3 $\pm 2.2$	18.2-24.5	20.9 $\pm 3.0$
18:2n-6	1.3- 2.0	1.6 $\pm 0.3$	3.6- 6.1	4.7 $\pm 1.2$	0.9- 1.2	1.0 $\pm 0.2$	3.9- 4.4	4.0 $\pm 0.3$
20:1n-9	1.0- 2.7	1.7 $\pm 0.9$	0.9- 1.9 (n=3)	1.4 $\pm 0.5$	0.5- 1.0	0.7 $\pm 0.3$	2.9- 5.1	4.1 $\pm 1.0$
20:4n-6	7.8- 9.8	8.6 $\pm 0.8$	6.8- 9.8	8.3 $\pm 1.6$	4.4- 7.0	5.3 $\pm 1.4$	4.1- 7.3	6.3 $\pm 1.5$
20:5n-3	8.6-10.6	9.3 $\pm 1.0$	5.0- 7.7	6.4 $\pm 1.5$	8.5- 8.9	8.8 $\pm 0.2$	2.9- 3.7	3.4 $\pm 0.3$
22:5n-3	1.4- 3.1	2.3 $\pm 0.7$	0.7- 1.3	0.8 $\pm 0.4$	2.0- 2.3	2.1 $\pm 0.2$	1.0- 1.2	1.1 $\pm 0.1$
22:6n-3	7.5-14.7	11.2 $\pm 3.0$	8.2-14.5	11.4 $\pm 2.6$	9.2-12.5	11.0 $\pm 1.7$	11.8-14.3	12.9 $\pm 1.1$
22:1n-9	0.4- 0.5 (n=3)	0.45 $\pm 0.007$	0.3- 0.7 (n=3)	0.49 $\pm 0.17$	0.2- 0.6	0.33 $\pm 0.16$	0.8- 0.9	0.86 $\pm 0.07$

are shown in table 1. Linoleic (18:2n-6) and arachidonic (20:4n-6) acids were significantly ( $P < 0.05$  and  $P < 0.025$ , respectively) higher and palmitic (16:0) acid significantly lower in the pre-moult rockhoppers ( $P < 0.025$ ) than the corresponding magellanics.

In both species of penguins moulting resulted in reduction in eicosapentaenoic (rockhoppers  $P < 0.025$ , magellanics  $P < 0.001$ ) and docosapentaenoic (22:5n-3) (rockhoppers  $P < 0.01$ , magellanics  $P < 0.001$ ) and an increase in linoleic acid percent (rockhoppers  $P < 0.005$ , magellanics  $P < 0.001$ ).

There was lower ( $P < 0.05$ ) stearic (18:0), and higher gadoleic (20:1n-9) ( $P < 0.005$ ) and erucic (22:1n-9) ( $P < 0.05$ ) acids in the post-moult magellanics compared to their pre-moult counterparts. The post-moult rockhoppers, however, had increased palmitic ( $P < 0.01$ ) and decreased palmitleic (16:1) ( $P < 0.05$ ) acids compared to the corresponding pre-moult birds.

Post-moult magellanics had significantly higher palmitoleic ( $P < 0.025$ ), gadoleic (20:1n-9) ( $P < 0.01$ ) and erucic (22:1n-9) ( $P < 0.05$ ) acids and significantly lower arachidonic ( $P < 0.05$ ), and eicosapentaenoic (20:5n-3) ( $P < 0.025$ ) acids compared to that of the post-moult rockhoppers.

## DISCUSSION

In both the rockhopper and magellanic penguins the major hepatic fatty acids were palmitic (16:0), stearic (18:0), oleic (18:1), arachidonic (20:4n-6), eicosapentaenoic (20:5n-3) and docosahexaenoic (22:6n-3). These findings are in general agreement with the reported fatty acid profiles for total body fat in wild Adelie (*Pygoscelis adeliae*) penguins [23] and dermal tissue in Emperor penguins (*Aptenodytes forsteri*) [24].

Pre-moult differences in hepatic fatty acids between rockhopper and magellanic penguins were likely to be due to differences in dietary habits. The rockhoppers including *Eudyptes crestatatus* feed opportunistically on squid, crustaceans, euphausiids, and small fish [11, 25, 26]. These foods would be rich in long chain n-3 fatty acids, with smaller amounts of the n-6 [27-30]. The magellanics' lower hepatic n-6 fatty acids imply a greater diet-

ary dependence on n-3 rich species. Diverse feeding ecologies have been reported for several penguin species [3, 31] and fatty acid compositions are known to be a reflection of both metabolism and diet [32-34]. Zar [24] suggested that the fatty acid differences between the adipose tissues of the Emperor (*Aptenodytes forsteri*) and of the Adelie (*Pygoscelis adeliae*) penguins were an effect of diet. In addition, Johnson and West [23] found that the proportions of fatty acids in Adelie penguin depot fat closely resembled the proportions of fatty acids in their normal diet of krill (*Euphausia sp.*).

Both penguin species have relatively high proportions of liver arachidonic acid (n-6) despite living in a n-3 fatty acid rich environment. This may indicate a physiological requirement in penguins for arachidonic acid similar to that in mammalian species (33). Diet selection patterns or rates of desaturation and elongation could account for the relatively high arachidonic acid composition. Similarly we previously reported [35] significant proportions of n-6 fatty acids in the liver phosphoglycerides of wild dolphins feeding in n-3 fatty acid-rich environments.

The post-moult disparity in hepatic fatty acids between the rockhoppers and magellanics was likely to be the result of species differences in the metabolism of lipids. It could also be that the penguins were at different stages of moulting, utilising nutrients differently. Moulting is characterised by three distinct phases, I and II representing essentially lipid catabolism, with >90% of energy expenditure stemming from lipids in phase II; in phase III any remaining lipid reserves, are catabolised and proteins are also utilised [36, 37]. The rockhoppers were perhaps in the final phase of moulting, utilising more proteins; and the magellanics in the earlier phases, because of their greater fatty acid mobilisation. These findings are in agreement with our earlier observations [5]; that the post-moult rockhoppers had significantly lower plasma albumin and globulin compared to their magellanic counterparts indicating that the rockhoppers were utilising higher amounts of protein.

Stearic acid (18:0) is quantitatively a major fatty acid in animal tissues, and is not preferentially oxidised as fuel in mammals [38]. The post-moult

decrease of stearic and concomitant increase in gadoleic (20:1n-9) and erucic (22:1n-9) acid percent in the magellanics but not in the rockhoppers, is therefore interesting. Increased elongation and desaturation of stearic acid in the magellanics to compensate for the relative post-moult loss in unsaturation, may explain these findings. Physiological adaptation to low environmental temperatures result in increase of unsaturation in the tissues of many species [39-42]. The post-moult drop in unsaturation in the magellanics, may have induced this adaptive elongation and desaturation mechanism. Preferential mobilisation and utilisation of the specific long chain n-3 fatty acids eicosapentaenoic and docosapentaenoic (22:5n-3) during moulting was a consistent biochemical finding in both the rockhopper and magellanics. Because of the dietary abundance of n-3 fatty acids in marine ecosystems, it appears that these penguins have evolved metabolic mechanisms to preferentially utilise these fatty acids.

There was a significant increase in the proportion of hepatic linoleic acid (18:2n-6) after moulting both in the magellanics and rockhoppers, together with a reduction in the long chain n-3 fatty acids, eicosapentaenoic (20:5n-3) and docosapentaenoic (22:5n-3). This n-6 and n-3 interaction is consistent with the findings of Gudbjarnason and Oskarsdottir [43] and Harbige *et al.* [22] in mammals. They found that increases in the proportion of long chain n-3 fatty acids was associated with a decrease in n-6 fatty acids, particularly linoleic acid. Linoleic acid is thought to have a role in the maintenance or formation of the epidermal water barrier [16]. It is conceivable that our observations of increased linoleic acid percent in the liver of both penguin species post-moult, may indicate specific mobilisation in relation to water barrier function during the vulnerable moulting period. Also there appears to be a differential sparing or conservation of the highly unsaturated docosahexaenoic acid (22:6n-3) and arachidonic acid in both species after moulting. Specific increases in membrane docosahexaenoic acid (22:6n-3) with cold adaptation have been reported [44, 45]; and may partly explain our findings as could the preferential retention of these fatty acids by hepatic cells. Conservation of docosahexaenoic acid, may also

have contributed to the post-moult decrease in the proportions of eicosapentaenoic and docosapentaenoic acid composition through chain elongation and or desaturation.

In conclusion we suggest that pre-moult hepatic fatty acid status of the rockhopper and magellanic penguins are mainly of dietary origin. Whereas, the post-moult values are a reflection of the stage of moult, and mobilisation and utilisation of lipids which appear to be both species dependent and species independent.

## REFERENCES

- 1 Moss, R., Watson, A., and Parr R. (1975) Maternal nutrition and breeding success in red grouse (*Lagopus lagopus scoticus*). *J. Animal. Ecol.*, **44**: 233-244.
- 2 Odum, E. P. (1965) Adipose tissue in migratory birds. In "Handbook of Physiology." Section 5, Adipose tissue. Ed. by A. E. Renold and G. F. Cahill American Physiological Society Washington D. C., pp. 37-43.
- 3 Heezik, Y. and Davis, L. (1990) Effects of food variability on growth rates, fledging sizes and reproductive success in the yellow-eyed penguin (*Megadyptes antipodes*). *Ibis*, **132**: 354-365.
- 4 Hornfeldt, B., Carlsson, B. G., Lofgren, O. and Eklund, U (1990) Effects of cyclic food supply on breeding performance in Tengmalm's owl (*Aegolius funereus*). *Can. J. Zool.*, **68**: 522-530.
- 5 Ghebremeskel, K., Williams, G., Keymer, I. F., Horsley, D. and Gardner D. A. (1989) Plasma chemistry of rockhopper (*Eudyptes crestatus*), magellanic (*Spheniscus magellanicus*) and gentoo (*Pygoscelis papua*) wild penguins in relation to moult. *Comp. Biochem. Physiol.*, **92A**: 43-47.
- 6 Moss, R. (1972) Food selection by red grouse (*Lagopus lagopus scoticus*) in relation to chemical composition. *J. Animal. Ecol.*, **41**: 411-428.
- 7 ChereI, Y. and Le Maho Y. (1988) Changes in body mass and plasma metabolites during short-term fasting in the king penguin. *The Condor*, **90**: 257-258.
- 8 Stonehouse, B. (1967) The general biology and thermal balances of penguins. *Adv. ecol. Res.*, **4**: 131-196.
- 9 Richdale, L. E. (1957) A population study of Penguins. Clarendon Press, Oxford, p. 95.
- 10 Sparks, J and Soper T. (1967) Penguins. by Newton Abbott: David & Charles, p. 101.
- 11 Strange, I. J. (1982) Breeding ecology of the rockhopper penguin (*Eudyptes crestatus*) in the Falkland Islands. *Le Gerfaut*, **72**: 137-188.
- 12 Williams, A. J., Siegfried, W. E., Burger, A. E. and

- Berruti A. (1977) Body composition and energy metabolism of moulting eudytid penguins. *Comp. Biochem. Physiol.*, **56A**: 27-30.
- 13 Gurr, M. I. and James A. T. (1980) Lipid Biochemistry: an introduction. Chapman and Hall, p. 111.
- 14 Crawford, M. A. (1983) Background to essential fatty acids and their prostanoid derivatives. *Br. Med. Bull.*, **39**: 210-213.
- 15 Martin, D. W., Mayes, P. A. and Rodwell V. W. (1983) Harper's Review of Biochemistry. 19th edition. Lange Medical Publications, pp. 188-247.
- 16 Wertz, P. W., Abraham, W., Cho, E. S. and Downing D. T. (1986) Linoleate-Rich O-Acylsphingolipids of mammalian epidermis: Structures and effects of essential fatty acid deficiency. *Prog. Lipid. Res.*, **25**: 383-389.
- 17 Hou, S. E., Mitra, A. K., White, S. H., Menon, G. K., Ghadially, R. and Elias P. M. (1991) Membrane structures in normal and essential fatty acid-deficient stratum corneum: Characterization by ruthenium tetroxide staining and X-ray diffraction. *J. Invest. Dermatol.*, **96**: 215-223.
- 18 Budowski, P., Leighfield, M. J. and Crawford, M. A. (1987) Nutritional encephalomalacia in the chick: an exposure of the vulnerable period for cerebellar development and the possible need for both n-6 and n-3 fatty acids. *Br. J. Nutr.*, **58**: 511-520.
- 19 Bazan, N. G. (1990) Supply of n-3 polyunsaturated fatty acids and their significance in the central nervous system. In "Nutrition and the brain." Vol. 8. Ed. by R. J. Wurtman and J. J. Wurtman Raven Press Ltd., NY. pp. 1-24.
- 20 Keymer, I. F. (1987) An investigation of rockhopper penguin (*Endiptes crestatus*) mortality in the Falkland Islands during the 1985-1986 breeding season. Falkland Island Foundation Project Report.
- 21 Folch, J. L. and Stanley, G. H. S. (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.*, **226**: 497-509.
- 22 Harbige, L. S., Ghebremeskel, K., Williams, G. and Summers, P. (1990) N-3 and N-6 phosphoglyceride fatty acids in relation to in vitro erythrocyte haemolysis induced by hydrogen peroxide in captive common marmosets (*Callithrix jacchus*). *Comp. Biochem. Physiol.*, **97B**: 1. 167-170.
- 23 Johnson, S. R. and West, G. C. (1973) Fat content, fatty acid composition and estimates of energy metabolism of adelic penguins (*Pygoscelis adeliae*) during the early breeding season fast. *Comp. Biochem. Physiol.*, **45B**: 709-719.
- 24 Zar, J. H. (1977) Fatty acid composition of emperor penguin (*Aptenodytes forsteri*) lipids. *Comp. Biochem. Physiol.*, **56B**: 109-112.
- 25 Brown, C. R. and Klages, N. T. (1987) Seasonal and annual variation in diets of Macaroni (*Eudyptes chrysolophus chrysolophus*) and Southern rockhopper (*E. chrysocome chrysocome*) penguins at sub-Antarctic Marion Island. *J. Zool., Lond.*, **212**: 7-28.
- 26 Croxall, J. P., Prince, P. A., Baird, A. and Ward, P. (1985) The diet of the southern rockhopper penguin *Eudyptes chrysocome chrysocome* at Beauchene Island, Falkland Islands. *J. Zool., Lond., (A)*, **206**: 485-496.
- 27 Crawford, M. A., Doyle, W., Drury, P., Ghebremeskel, K., Harbige, L., Leyton, J. and Williams, G. (1989) The food chain for n-6 and n-3 fatty acids with special reference to animal products. In "Dietary n-3 and n-6 Fatty Acids biological effects and nutritional essentiality." NATO ASI Series A, Vol 171. Ed. by C. Galli. and A. P. Simopoulos Plenum Press, pp. 5-19.
- 28 Ackman, R. G. (1982) Fatty acid composition of fish oils. In "Nutritional evaluation of long-chain fatty acids in fish oil." Ed. by S. M. Barlow and M. E. Stansby Academic Press, pp. 25-88.
- 29 Sidhu, G. S., Montgomery, W. A., Holloway, G. L., Johnson, A. R. and Walker D. M. (1970) Biochemical composition and nutritive value of krill (*Euphausia superba*). *J. Sci. Fd. Agric.*, **21**: 293-296.
- 30 Clarke, A. (1980) The biochemical composition of krill *Euphausia superba*, from South Georgia. *J. exp. mar. Biol. Ecol.*, **43**: 221-236.
- 31 Lishman G. S. (1985) The food and feeding ecology of adelic penguins (*Pygoscelis adeliae*) and Chinstrap penguins (*P. antarctica*) at Signy Island, South Orkney Islands. *J. Zool., Lond., (A)*, **205**: 245-263.
- 32 Tinco, J. (1982) Dietary requirements and functions of alpha-linolenic acid in animals. *Prog. Lipid. Res.*, **21**: 1-45.
- 33 Sanders, T. A. B. (1988) Essential and Trans-fatty acids in nutrition. *Nutrition Research Reviews*, **1**: 57-78.
- 34 Stubbs, C. D. and Smith, A. D. (1984) The modification of mammalian membrane polyunsaturated fatty acid composition in relation to membrane fluidity and function. *Biochim. Biophys. Acta.*, **779**: 89-137.
- 35 Williams, G. and Crawford M. A. (1987) Comparison of the fatty acid component in structural lipids from dolphins, zebra and giraffe: possible evolutionary implications. *J. Zool., Lond.*, **213**: 673-684.
- 36 Cherel, Y., Robin J-P. and Le Maho Y. (1988) Physiology and biochemistry of long-term fasting in birds. *Can. J. Zool.*, **66**: 159-166.
- 37 Robin, J-P., Frain, M., Sardet, C., Groscolas, R. and Le Maho Y. (1988) Protein and lipid utilization during long-term fasting in emperor penguins. *Am. J. Physiol.*, **254**: R61-R68.
- 38 Leyton, J., Drury, P. J. and Crawford, M. A. (1987)

- In vivo incorporation of labeled fatty acids in rat liver lipids after oral administration. *Lipids*, **228**: 553-558.
- 39 Williams, D. D. and Platner W. S. (1967) Cold-induced changes in fatty acids of the rat and hamster. *Am. J. Physiol.*, **212**(1): 167-172.
- 40 Aloia, R. C. and Pengelley, E. T. (1979) Lipid composition of cellular membranes of hibernating mammals. In "Chemical Zoology." Vol XI Mammalia. Ed. by M. Florkin and B. T. Scheer Academic Press, pp. 1-47.
- 41 Cowey, C. B. (1988) The Nutrition of fish: The developing scene. *Nutrition Research Reviews*, **1**: 255-280.
- 42 Houslay, M. D., and Stanley, K. K. (1983) Dynamics of Biological Membranes. John Wiley & Sons, pp. 138-141.
- 43 Gudbjarnason, S. and Oskarsdottir, G. (1977) Modification of fatty acid composition of rat heart lipids by feeding cod liver oil. *Biochim. Biophys. Acta.*, **487**: 10-15.
- 44 Miller, N. G. A., Hill, M. W. and Smith M. W. (1976) Positional and species analysis of membrane phospholipids extracted from goldfish adapted to different environmental temperatures. *Biochim. Biophys. Acta.*, **55**: 644-654.
- 45 Hazel, J. R. and Zerba, E. (1986) Adaption of biological membranes to temperature: molecular species compositions of phosphatidyl choline and phosphatidyl ethanolamine in mitochondrial and microsomal membranes of liver from thermally acclimated rainbow trout. *J. Comp. Physiol.* **156B**: 665-674.